

# **Classifying and Prioritizing Watersheds for Protection and Restoration: An Approach for the Mid-Atlantic Highlands Area**

## **Sponsor:**

U.S. Environmental Protection Agency, Science to Achieve Results (STAR) Program, EPA-STAR Grant #R-830593

## **Project Officer:**

Iris Goodman, US EPA, ORD, National Center for Environmental Research

## **Project Co-Directors:**

Robert P. Brooks (Penn State) and George D. Constantz (Canaan Valley Institute)

## **Institutions:**

The Pennsylvania State University, University Park, PA  
Cooperative Wetlands Center  
Department of Geography  
School of Forest Resources  
Center for Statistical Ecology and Environmental Statistics  
Penn State Institutes of Energy and the Environment  
Canaan Valley Institute, Davis, WV and Clearfield, PA

## **Report Authors:**

Bronson Griscom, Mary McKenney-Easterling, Wayne L. Myers, Abby McQueen, Joseph A. Bishop, Gian Rocco, Janie French, Kristen C. Hychka, Alynne Bayard, Timothy O'Connell, G.P. Patil, Charles Taillie (deceased), George D. Constantz and Robert P. Brooks

## SYNOPSIS

Natural resource managers need to be able to put watersheds into categories for several reasons, including identifying reference conditions, understanding types of environmental degradation, designing monitoring studies, and narrowing restoration options. In this project, we explored ways to develop and apply a hierarchical, geographically-independent classification of watersheds based on pre-existing environmental data. Our work was focused on the Mid-Atlantic Highlands Area, but has applications elsewhere.

Our approach in this project was to first construct a classification system that characterizes watersheds based on their inherent natural features (climate, soils, topography and hydrology). This resulted in the delineation of nine distinct clusters of watersheds for the region. We then developed an approach to prioritization that classifies watersheds according to human disturbance (primarily expressed through land cover) and their susceptibility to impairment from a variety of stressors, including land use, acidification, impervious cover, and nutrients. Based on our research products and conversations with potential users, we discovered that the best way to “prioritize” watersheds was use an interactive process. We produced maps of our nine cluster groups showing their spatial distribution in the region. These maps can be used by managers to recognize where and how their specific watersheds of concern fit into a larger landscape context. We produced narrative descriptions of the nine watershed clusters that relate their inherent characteristics to their vulnerability of being impacted by a suite of stressors. This approach is preferred to one where researchers impose a prioritization scheme on potential users. This method also allows users to incorporate other information and data that address the invariably unique issues at hand for that single watershed or group of watersheds. Given that we delineated over 2,800 small watersheds in the study area, this seemed to be the most appropriate approach.

We tested the efficacy of our classification system across watersheds in the state of West Virginia, where the best synoptic measure of ecological integrity was available. Using measures of land use in the watershed and riparian corridor, and stream biological integrity, we found that that the nine clusters of watersheds were sufficiently differentiated that they showed variations in vulnerability to human-generated stressors. Vulnerability (ecological resistance) tended to have an inverse relationship with likelihood of land use impacts: high vulnerability watershed classes tended to have relatively low land use impacts, while low vulnerability watershed classes tended to have relatively high land use impacts.

Our recommendations to potential users are to locate watersheds of interest, determine their cluster membership, and then consider their vulnerability to expected stressors. To make this process convenient for users, we produced a Watershed Characterization and Prioritization Tool that helps users locate graphically watersheds of interest and obtain both characterization data and an initial assessment of their vulnerability. By understanding the inherent characteristics that define each cluster, and considering the watershed’s probable response to specific stressors, one can conceptually locate a watershed in the two dimensional space, portrayed in Figure S1 below. When it is necessary to prioritize among multiple watersheds, then the relative location of a watershed in that space will suggest which is most vulnerable and in need of attention first.

We believe this approach is simple for, adaptable to, and useful by managers, because it combines the best available information from scientific investigations with the knowledge and intentions of local stakeholders. Whether comparing among watersheds or varying condition within the same cluster type or across cluster types, this approach should generate a relevant list of prioritized watersheds. The contemplative process used to locate multiple watersheds in this conceptual space should be most helpful in deciding upon a course of action with regard to prioritizing watershed protection and restoration.





Figure S1. Conceptual framework for prioritizing watersheds for protection and restoration, according to their location in two-dimensional space defined by axes of disturbance and vulnerability to impairment.

Figure S2. Schematic flow chart of watershed characterization and prioritization analyses.



## TABLE OF CONTENTS

SYNOPSIS	i
ACKNOWLEDGEMENTS	v
INTRODUCTION	1
Rationale and Context	1
Objectives and Approach	2
Study Area	3
CHARACTERIZATION BASED ON INHERENT ATTRIBUTES OF WATERSHEDS	6
Dataset Development and Exploratory Analyses	6
Development of Candidate Classifications	8
Naming of the Nine Clusters	8
CHARACTERIZATION OF STRESSORS	11
Land Cover Stressors	12
In-stream Stressors	14
STRESSOR CLASS IDENTIFICATION AND CONDITION RANKING	17
VULNERABILITY OF WATERSHEDS TO HUMAN IMPACTS IN WEST VIRGINIA	21
Evaluation of Classification Systems	22
Comparison of Vulnerability among Inherent Classes	24
DISCUSSION & CONCLUSIONS	28
Vulnerability of Watershed Classes	29
Watershed Classification: Description of classes and vulnerability	32
Recommendations for Applying Watershed Classification Approach	38
STAKEHOLDER COMMUNICATION AND WATERSHED TOOL	40
REFERENCES & DATASETS	43
APPENDICES	
Appendix A. Full set of candidate metrics computed for the Mid-Atlantic Highlands Area	
Appendix B. List of manuscripts and other files on CD.	
Appendix C. Summary of meetings with stakeholders.	

## ACKNOWLEDGMENTS

This project would not have been possible without the collaborative teamwork of all participants from Penn State and CVI. In addition, we thank the various stakeholders who agreed to comment on earlier drafts of our work. This research has been supported by a grant from the U.S. Environmental Protection Agency's Science to Achieve Results (STAR) Grants Program, U.S. EPA Agreement CR-83059301. Although the research described in this report has funded wholly or in part by the United State Environmental Protection Agency, it has not been subjected to the Agency's required peer and policy review and, therefore, does not reflect the view of the Agency and no official endorsement should be inferred.

Robert P. Brooks and George Constantz, Co-Directors, April 2007

The recommended citation for this report is:

Griscom, B., M. McKenney-Easterling, W. L. Myers, A. McQueen, J. A. Bishop, G. Rocco, J. French, K. C. Hychka, A. Bayard, T. O'Connell, G.P. Patil, C. Taillie (deceased), G. D. Constantz and R. P. Brooks. 2007. Classifying and prioritizing watersheds for protection and restoration: an approach for the Mid-Atlantic Highlands Area. Final Report to the U.S. Environmental Protection Agency STAR Program, Agreement CR-83059301, Washington, DC. Prepared by Pennsylvania State University, University Park, PA and Canaan Valley Institute, Thomas, WV. 50pp.+appendices & CD.